

DOMAIN	Participant Demographics	Intervention Characteristics	Pre/Within/Post/Follow-Up Treatment measures  (please provide for all time points available at the individual item level)
<b>VARIABLES</b>	Age  Gender/sex  Race/Ethnicity  Socioeconomic status  Level of education  Method of recruitment into study  Marital Status  Acculturation status (if available)  Parental Status (if available)	<b>PTSD + SUD Intervention</b> <ul style="list-style-type: none"> <li>• Intervention used (name, citation)</li> <li>• How many sessions have an explicit focus on SUD (if any)?</li> <li>• How many sessions have an explicit focus on PTSD (if any)?</li> <li>• Intervention frequency (# sessions/week)</li> <li>• Intervention duration (# of weeks)</li> <li>• Qualifications of therapist(s)</li> <li>• Location of sessions</li> <li>• How did you address fidelity?</li> <li>• Number of sessions attended</li> </ul> <b>Comparator</b> <ul style="list-style-type: none"> <li>• Intervention used (name, citation)</li> <li>• How many sessions have an explicit focus on SUD (if any)?</li> <li>• How many sessions have an explicit focus on PTSD (if any)?</li> <li>• Intervention frequency (# sessions/week)</li> <li>• Intervention duration (# of weeks)</li> <li>• Qualifications of therapist(s)</li> <li>• Location of sessions</li> </ul>	<b>PTSD (please provide data at symptom/item level)</b> <ul style="list-style-type: none"> <li>• Clinical diagnostic tool used to diagnose PTSD</li> <li>• Clinical severity and frequency rating scales for PTSD (clinician and self-report at individual item level)</li> <li>• Post-traumatic cognitions (at individual item level)</li> <li>• Trauma during treatment (i.e., domestic violence; at individual item level)</li> <li>• Subjective units of distress scores</li> </ul> <b>Substance Use Disorder (please provide data at symptom/item level)</b> <ul style="list-style-type: none"> <li>• Clinical diagnostic tool used to diagnose SUD</li> <li>• Clinical severity rating scales for SUD (clinician and self-report at individual item level)</li> <li>• Frequency and quantity of daily substance use (all classes available; individual item level)</li> <li>• Days of each substances used</li> <li>• Primary substance used</li> <li>• SUD physiological measures (e.g., saliva, urine analysis, breathalyzer)</li> <li>• Craving measures (individual item level)</li> </ul> <b>Other Treatments</b> <ul style="list-style-type: none"> <li>• Use of psychotropic medications</li> <li>• Addiction-related medication</li> <li>• Specify Medications, dose taken, and dose available</li> </ul>

	<p>Military vs. Civilian Status</p> <p>If Military: active duty or veteran?</p> <p><b>Trauma characteristics</b></p> <p>Trauma types (child v. adult onset, total number of traumas, civilian or military, age of onset of trauma exposure domestic violence)</p> <p><b>Other diagnoses</b></p> <p>Non-psychiatric comorbidities</p>	<ul style="list-style-type: none"> <li>• How did you address fidelity?</li> <li>• Number of sessions attended</li> <li>•</li> </ul>	<ul style="list-style-type: none"> <li>• Treatment Services Review (individual item level)</li> </ul> <p><b>Other diagnoses and problems</b></p> <ul style="list-style-type: none"> <li>• Clinical diagnostic tool used for other mental health disorders (non-PTSD)</li> <li>• Other lifetime and current mental health disorders (individual item level)</li> </ul> <p>Functional rating scales (at individual item level):</p> <ul style="list-style-type: none"> <li>• Depression severity, anxiety severity</li> <li>• Global, social, functional impairment (e.g., Brief Symptom Inventory)</li> </ul> <p>Emotion/emotion dysregulation indices (at individual item level):</p> <ul style="list-style-type: none"> <li>• Emotion (e.g., PANAS)</li> <li>• Emotion dysregulation (e.g., DERS)</li> </ul> <p>Suicide/self-harm ideation/behavior/urges (at individual item level)</p> <p>Sleep problems/severity measures (at individual item level)</p> <p>Serious/Adverse Events</p> <p>Therapeutic alliance (at individual item level)</p>
--	--	---	---